

HOMOLOGICAL STABILITY FOR FAMILIES OF COXETER GROUPS

RICHARD HEPWORTH


ABSTRACT. We prove that certain families of Coxeter groups and inclusions $W_1 \hookrightarrow W_2 \hookrightarrow \dots$ satisfy homological stability, meaning that in each degree the homology $H_*(BW_n)$ is eventually independent of n . This gives a uniform treatment of homological stability for the families of Coxeter groups of type A_n , B_n and D_n , recovering existing results in the first two cases, and giving a new result in the third. The key step in our proof is to show that a certain simplicial complex with W_n -action is highly connected. To do this we show that the barycentric subdivision is an instance of the ‘basic construction’, and then use Davis’s description of the basic construction as an increasing union of chambers to deduce the required connectivity.

1. INTRODUCTION

A family of groups $G_1 \hookrightarrow G_2 \hookrightarrow G_3 \hookrightarrow \dots$ is said to satisfy *homological stability* if the induced maps $H_i(BG_{n-1}) \rightarrow H_i(BG_n)$ are isomorphisms when n is sufficiently large relative to i . Homological stability is known for many families of groups, including symmetric groups [13], general linear groups [14], mapping class groups of surfaces [7] and 3-manifolds [11], diffeomorphism groups of highly connected manifolds [6], and automorphism groups of free groups [8], [10]. *Coxeter groups* are abstract reflection groups, appearing in many areas of mathematics, such as root systems and Lie theory, geometric group theory, and combinatorics. See [2], [5], and [1] for introductions to Coxeter groups from each of these three viewpoints. In this paper we will show that homological stability holds for certain families of Coxeter groups.

Recall that a *Coxeter matrix* on a set S is an $S \times S$ symmetric matrix M , with values in $\mathbb{N} \cup \{\infty\}$, satisfying $m_{st} = 1$ if $s = t$ and $m_{st} \geq 2$ otherwise. The corresponding *Coxeter group* is the group generated by the elements of S , subject to the relations $(st)^{m_{st}} = e$ for $s, t \in S$. (When $m_{st} = \infty$ no relation is imposed.) It is common to represent a Coxeter matrix by the equivalent *Coxeter diagram*. This is the graph with vertices S and edges $\{s, t\}$ for $m_{st} \geq 3$. The edge $\{s, t\}$ is labelled m_{st} if $m_{st} \geq 4$.

Now consider a sequence of finite Coxeter diagrams $(\Gamma_n)_{n \geq 1}$ of the form



2010 *Mathematics Subject Classification.* 20F55, 20J06.

Key words and phrases. Homological stability, Coxeter groups.

where every diagram has a preferred vertex, and each diagram is obtained from its predecessor by attaching a new preferred vertex to the old one by an unlabelled edge. Writing W_n for the Coxeter group determined by Γ_n , the inclusion $\Gamma_{n-1} \hookrightarrow \Gamma_n$ induces an inclusion $W_{n-1} \hookrightarrow W_n$, and our main result states that the family


$$W_1 \hookrightarrow W_2 \hookrightarrow W_3 \hookrightarrow W_4 \hookrightarrow \dots$$

satisfies homological stability.

Main Theorem. *The map $H_*(BW_{n-1}) \rightarrow H_*(BW_n)$ is an isomorphism in degrees $2* \leq n$. Here homology is taken with arbitrary constant coefficients.*

Observe that while the diagrams Γ_n are assumed to be finite, it is not necessary for the groups W_n to be finite.

Homological stability for Coxeter groups of type A_n , B_n and D_n . The main theorem gives a uniform treatment of homological stability for the families of Coxeter groups of type A_n , B_n and D_n . Recall that these are the Coxeter groups corresponding to the following diagrams, in which n always denotes the total number of vertices.



These families have an important place in the theory of Coxeter groups, since the classification of finite Coxeter groups states that a finite irreducible Coxeter group has type A_n , B_n or D_n , or is dihedral, or is one of six exceptional examples. (See Appendix C of [5].) The sequences $(A_n)_{n \geq 1}$, $(B_{n+1})_{n \geq 1}$ and $(D_{n+2})_{n \geq 1}$ all have the form $(\Gamma_n)_{n \geq 1}$ described above, with the rightmost vertex taken as the preferred vertex, and therefore we may apply the main theorem to each one.

For the sequence of diagrams $(A_n)_{n \geq 1}$, the corresponding sequence of Coxeter groups is

$$\Sigma_2 \hookrightarrow \Sigma_3 \hookrightarrow \Sigma_4 \hookrightarrow \Sigma_5 \hookrightarrow \dots$$

where Σ_n is the symmetric group on n letters and the inclusions are given by extending permutations by the identity. Applying the main theorem, we recover the following classical result.

Corollary (Nakaoka [13]). *The map $H_*(B\Sigma_n) \rightarrow H_*(B\Sigma_{n+1})$ is an isomorphism in degrees $2* \leq n$.*

For the sequence of diagrams $(B_{n+1})_{n \geq 1}$, the corresponding sequence of Coxeter groups

$$C_2 \wr \Sigma_2 \hookrightarrow C_2 \wr \Sigma_3 \hookrightarrow C_2 \wr \Sigma_4 \hookrightarrow C_2 \wr \Sigma_5 \hookrightarrow \dots$$

consists of the wreath products of the symmetric groups with the group C_2 of order 2, and the inclusions are again given by extending permutations by the identity. Applying the main theorem gives the following special case of Hatcher and Wahl's result on homological stability for wreath products. (See Proposition 1.6 of [11] and the discussion that follows it.) It also follows from Randal-Williams's result on homological stability for unordered configuration spaces. (See Theorem A of [15] with $M = \mathbb{R}^\infty$ and $X = BC_2$.)

Corollary (Hatcher-Wahl [11]). *The map $H_*(B(C_2 \wr \Sigma_n)) \rightarrow H_*(B(C_2 \wr \Sigma_{n+1}))$ is an isomorphism in degrees $2* \leq n$.*

For the sequence of diagrams $(D_{n+2})_{n \geq 1}$, the corresponding sequence of Coxeter groups is


$$H_3 \hookrightarrow H_4 \hookrightarrow H_5 \hookrightarrow H_6 \hookrightarrow \dots$$

where H_n denotes the kernel of the homomorphism $C_2 \wr \Sigma_n \rightarrow C_2$ that takes the sum of the C_2 -components. The main theorem gives the following result, which we believe to be new.

Corollary. *Let H_n denote the Coxeter group of type D_n . Then the inclusion $H_{n+1} \hookrightarrow H_{n+2}$ induces an isomorphism $H_*(BH_{n+1}) \rightarrow H_*(BH_{n+2})$ in degrees $2* \leq n$.*

The concrete descriptions of the groups of type A_n , B_n and D_n that we used here can all be found in section 6.7 of [5].

Families of hyperbolic Coxeter groups. The main theorem applies to interesting families besides those of type A_n , B_n and D_n already considered. For example, if we fix an integer $m \geq 7$, then the main theorem shows that homological stability holds for the family of Coxeter groups associated to the sequence of diagrams $(\Gamma_n)_{n \geq 1}$



in which Γ_n has a total of $(n + 1)$ vertices, the rightmost one preferred. This family has the feature that the first group is finite, while the rest are all infinite hyperbolic. (Hyperbolicity is verified using Moussong's condition. See Corollary 12.6.3 of [5].) It is not difficult to create other sequences of infinite hyperbolic groups to which the main theorem applies.

On the other hand, these examples show that it is possible for the first group in one of our families to be finite while the rest are infinite. Similarly, it is possible for the first group to be infinite hyperbolic while the rest are not, for example if $\Gamma_1 = \bullet \xrightarrow{6} \bullet \xrightarrow{6} \bullet$ with any preferred vertex. (The claims about hyperbolicity again follow from Moussong's condition.)

Overview of the proof. The proof of the main theorem proceeds as follows.

First, we construct a simplicial complex $\mathcal{C}^n$ with an action of W_n , and prove that it is weakly Cohen-Macaulay of dimension n , meaning that it is $(n - 1)$ -connected in a certain 'homogeneous' way.

Second, we form the semisimplicial set $\mathcal{D}^n$ whose simplices are simplices of $\mathcal{C}^n$ with an ordering of their vertices. We show that W_n acts transitively on the simplices in each dimension, with stabilisers the subgroups $W_m \subset W_n$ for $m < n$. From the weakly Cohen-Macaulay property of $\mathcal{C}^n$ we deduce that the geometric realisation $\|\mathcal{D}^n\|$ is $(n - 1)$ -connected.

Third, we study the Borel construction $EW_n \times_{W_n} \|\mathcal{D}^n\|$. From properties of $\mathcal{D}^n$ we deduce that the homology of the Borel construction matches that of BW_n in a range of degrees, and that it can be computed by a spectral sequence whose

E^1 -term consists of the homology groups $H_*(BW_m)$ for $m < n$. An argument involving this spectral sequence completes the proof.

Let us explain in more detail how we show that $\mathcal{C}^n$ is $(n-1)$ -connected, since this is by far the hardest step in the proof. We make use of the ‘basic construction’, a technique from the topology of Coxeter groups (see chapters 5 and 8 of Davis’s book [5]). The basic construction takes a Coxeter system (W, S) and a ‘mirrored space’ X over S , and produces a topological space $\mathcal{U}(W, X)$ with W -action. Now $\mathcal{U}(W, X)$ can be expressed as an ‘increasing union of chambers’, i.e. copies of X , each copy of X being attached to the preceding ones in a controlled way. To prove that $\mathcal{C}^n$ is $(n-1)$ -connected, we show that the realisation of the barycentric subdivision of $\mathcal{C}^n$ has the form $\mathcal{U}(W_n, |\Delta|)$, where $|\Delta|$ is a topological n -simplex. By studying the attachments in the increasing union of chambers, we are able to deduce the required connectivity.

Outline of the paper. In section 2 we establish some notation, and in section 3 we prove some elementary algebraic facts about the groups W_n . In section 4 we define the simplicial complex $\mathcal{C}^n$. Then we examine $\mathcal{C}^n$ in detail: in section 5 we study the links of its simplices, in section 6 we study its action by W_n , and in section 7 we show that it is $(n-1)$ -connected. Then in section 8 we study the semisimplicial set of ordered simplices in $\mathcal{C}^n$. The proof of the main theorem is completed in section 9.

Acknowledgments. My thanks to Jarek Kędra, Ian Leary and Oscar Randal-Williams for helpful conversations as this work was being carried out.


2. NOTATION

Definition 1. Let $(\Gamma_n)_{n \geq 1}$ be a sequence of the kind described in the introduction. We extend this sequence to the left by two terms as follows. Define Γ_0 to be the diagram obtained from Γ_1 by deleting the preferred vertex. And define Γ_{-1} to be the diagram obtained from Γ_1 by deleting the preferred vertex and all vertices that shared an edge with it.

Example 2 (Coxeter groups of type A_n , B_n and D_n). For the sequence $(A_n)_{n \geq 1}$, the diagrams A_0 and A_{-1} are both empty. For $(B_{n+1})_{n \geq 1}$, the diagram B_{0+1} consists of a single vertex and B_{-1+1} is empty. And for $(D_{n+2})_{n \geq 1}$, the diagram D_{0+2} consists of two vertices with no edge and D_{-1+2} is empty.

Definition 3. Let $(\Gamma_n)_{n \geq 1}$ be a sequence of the kind described in the introduction, and let $(\Gamma_n)_{n \geq -1}$ be the extension described above. Then for $n \geq -1$ we define S_n to be the set of vertices of Γ_n , and we define W_n to be the Coxeter group associated to Γ_n . Thus (W_n, S_n) is a Coxeter system.

Definition 4. For $n \geq 1$ we define $s_n \in S_n$ to be the preferred vertex of Γ_n .



Thus $S_n = S_0 \cup \{s_1, \dots, s_n\}$, and the parabolic subgroup of W_n generated by $s_1, \dots, s_n$ is an isomorphic copy of Σ_{n+1} , with s_i acting as the transposition of i and $(i+1)$.

3. ALGEBRAIC PRELIMINARIES

From this point onwards, unless stated otherwise we fix the sequence $(\Gamma_n)_{n \geq 1}$ and the integer $n \geq 1$.

In several places we will consider the symbol $s_i \cdots s_n W_{n-1}$ for i in the range $1 \leq i \leq n+1$. In the case $i = n+1$ we take it to mean W_{n-1} .

Proposition 5. *Let i lie in the range $1 \leq i \leq n$. Then left multiplication by the element s_i fixes the set*

$$\{s_1 \cdots s_n W_{n-1}, s_2 \cdots s_n W_{n-1}, \dots, s_n W_{n-1}, W_{n-1}\}.$$

It acts on the set by transposing $s_i \cdots s_n W_{n-1}$ and $s_{i+1} \cdots s_n W_{n-1}$, and fixing the remaining elements.

Proof. The identities

$$\begin{aligned} s_i(s_j \cdots s_n) &= (s_j \cdots s_n)s_i \text{ for } i < j-1 \\ s_i(s_{i+1} \cdots s_n) &= s_i \cdots s_n \\ s_i(s_i \cdots s_n) &= s_{i+1} \cdots s_n \\ s_i(s_j \cdots s_n) &= (s_j \cdots s_n)s_{i-1} \text{ for } i > j \end{aligned}$$

are simple to verify, and the claim follows immediately. $\square$

Proposition 6. $W_{i-1} \cap (s_i \cdots s_n W_{n-1} s_n \cdots s_i) = W_{i-2}$ for $1 \leq i \leq n$.

Proof. The word $s_i \cdots s_n$ is (W_{i-1}, W_{n-1}) -reduced, meaning that it is reduced and has no reduced expression beginning with a generator of W_{i-1} or ending with a generator of W_{n-1} . A result of Kilmoyer, Solomon and Tits (see Lemma 2 of [16] and the remarks that precede it) then shows that $W_{i-1} \cap s_i \cdots s_n W_{n-1} s_n \cdots s_i$ is the subgroup generated by $T = S_{i-1} \cap (s_i \cdots s_n S_{n-1} s_n \cdots s_i)$. So it will be enough to show that $T = S_{i-2}$. It is immediate that $S_{i-2} \subset T$, so suppose that $t \in T \setminus S_{i-2}$. Thus $t \in S_{i-1} \setminus S_{i-2}$ and $s_n \cdots s_i t s_i \cdots s_n \in S_{n-1}$. By the first condition we have $m(s_i, t) \geq 3$. By the second condition the word $s_n \cdots s_i t s_i \cdots s_n$ is not reduced, so that we must be able to apply an M-move to it (see section 3.4 of [5]), and this is only possible if $m(s_i, t)$ is exactly 3. But in this case $s_n \cdots s_i t s_i \cdots s_n$ is already reduced, contradicting the first condition. $\square$

Proposition 7. *Let i lie in the range $1 \leq i \leq n$. If $\sigma, \tau \in W_n$ satisfy*

$$\sigma s_j \cdots s_n W_{n-1} = \tau s_j \cdots s_n W_{n-1} \text{ for } j = i, \dots, n+1$$

then $\sigma^{-1}\tau \in W_{i-2}$.

Proof. The proposition is equivalent to the claim that

$$W_{n-1} \cap (s_n W_{n-1} s_n) \cap \cdots \cap (s_i \cdots s_n W_{n-1} s_n \cdots s_i) = W_{i-2},$$

which is proved by downward induction on i . The initial case $i = n+1$ is immediate, and the induction step follows from Proposition 6. $\square$

Proposition 8. *For $c \in W_n$ the cosets*

$$c(s_1 \cdots s_n)W_{n-1}, \dots, cs_n W_{n-1}, cW_{n-1}$$

are pairwise distinct.

Proof. If $cs_j \cdots s_n W_{n-1} = cs_k \cdots s_n W_{n-1}$ with $j < k$, then $(s_n \cdots s_j)(s_k \cdots s_n) \in W_{n-1}$. But $(s_n \cdots s_j)(s_k \cdots s_n) = (s_{k-1} \cdots s_n \cdots s_{k-1})(s_{k-2} \cdots s_j)$, implying that $s_n \in W_{n-1}$, which is a contradiction. $\square$

4. THE SIMPLICIAL COMPLEX $\mathcal{C}^n$

Now we introduce the simplicial complex that will be central to our proof of the main theorem, and we prove that it is weakly Cohen-Macaulay of dimension n . The proof relies on propositions that will be established in the following three sections.

Definition 9 (The simplicial complex $\mathcal{C}^n$). Given $n \geq 0$ we let $\mathcal{C}^n$ denote the n -dimensional abstract simplicial complex with vertex set W_n/W_{n-1} and with k -simplices given by the subsets

$$C = \{c(s_{n-k+1} \cdots s_n)W_{n-1}, \dots, cs_n W_{n-1}, cW_{n-1}\}$$

for $0 \leq k \leq n$ and $c \in W_n$. (Proposition 8 shows that C does indeed have cardinality $(k+1)$.) In this situation we call c a *lift* of the simplex C .

Remark 10. We chose the name “lift” in the previous definition to emphasise the formal similarity with the concept of the same name that appears in Definition 2.1 of Wahl’s paper [17].

A given simplex can have many lifts. Choosing a lift for a simplex induces an ordering of its vertices, and all orderings occur in this way. For if c lifts a k -simplex C then so does $cs_{n-k+i+1}$, and the induced orderings differ by transposition of the i -th and $(i+1)$ -st vertices (see Proposition 5). This makes it simple to verify that $\mathcal{C}^n$ is indeed a simplicial complex, for if C is a simplex of $\mathcal{C}^n$ and $D \subset C$ is a nonempty subset, then we may choose a lift c of C such that D is a terminal segment in the induced ordering. Then c is also a lift of D .

The natural action of W_n on W_n/W_{n-1} extends to an action on $\mathcal{C}^n$. For if C is a simplex of $\mathcal{C}^n$ with lift c , and if $w \in W_n$, then wC is a simplex of $\mathcal{C}^n$ with lift wc .

We now give a concrete description of $\mathcal{C}^n$ for the families of Coxeter groups of type A_n , B_n and D_n that were discussed in the introduction. Again, see section 6.7 of [5] for the concrete descriptions of these groups.

Example 11 (The family A_n). Let the sequence of diagrams $(\Gamma_n)_{n \geq 1}$ be $(A_n)_{n \geq 1}$, so that $W_n = \Sigma_{n+1}$ is the symmetric group on $(n+1)$ letters. Then $\mathcal{C}^n$ is the n -dimensional simplex Δ^n with the action of Σ_{n+1} that permutes the vertices. For the vertex set of $\mathcal{C}^n$ is Σ_{n+1}/Σ_n , which is isomorphic to $\{1, \dots, n+1\}$ via the map that sends $\sigma\Sigma_n$ to $\sigma(n+1)$. Under this isomorphism, an element $\sigma \in \Sigma_{n+1}$ is a lift of the k -simplex

$$C = \{\sigma(n-k+1), \dots, \sigma(n+1)\},$$

and every subset of $\{1, \dots, n+1\}$ arises in this way.

Example 12 (The family B_n). Let the sequence of diagrams $(\Gamma_n)_{n \geq 1}$ be given by $(B_{n+1})_{n \geq 1}$, so that $W_n = C_2 \wr \Sigma_{n+1}$. In this case $\mathcal{C}^n$ is isomorphic to the *hyperoctahedron of dimension n* , which is the simplicial complex whose vertex set is $\{\pm 1, \dots, \pm(n+1)\}$ and whose simplices are the subsets containing at

most one element from each pair $\{i, -i\}$. In particular, its realisation is homeomorphic to the n -sphere. The action of $C_2 \wr \Sigma_{n+1}$ on the hyperoctahedron is the one in which Σ_{n+1} permutes the pairs $\pm i$ while preserving their signs, and in which the i -th copy of C_2 transposes i and $-i$. To obtain this description, observe that the vertex set of $\mathcal{C}^n$ is $(C_2 \wr \Sigma_{n+1})/(C_2 \wr \Sigma_n)$, which is isomorphic to $\{\pm 1, \dots, \pm(n+1)\}$ via the map that sends the coset of $((\epsilon_1, \dots, \epsilon_{n+1}), \sigma)$ to $\epsilon_{n+1}\sigma(n+1)$. (Here we are taking $C_2 = \{\pm 1\}$.) And under this isomorphism an element $((\epsilon_1, \dots, \epsilon_{n+1}), \sigma)$ lifts the k -simplex

$$C = \{\epsilon_{n-k+1}\sigma(n-k+1), \dots, \epsilon_{n+1}\sigma(n+1)\},$$

so that a subset of $\{\pm 1, \dots, \pm(n+1)\}$ spans a simplex of $\mathcal{C}^n$ if and only if it does not contain any element and its negative.

Example 13 (The family D_n). Let the sequence of diagrams $(\Gamma_n)_{n \geq 1}$ be given by $(D_{n+2})_{n \geq 1}$, so that $W_n = H_{n+2}$ is the kernel of the homomorphism $C_2 \wr \Sigma_{n+2} \rightarrow C_2$ that takes the sum of the C_2 -components. In this case $\mathcal{C}^n$ is the n -skeleton of the $(n+1)$ -dimensional hyperoctahedron, with the action inherited from the action of $C_2 \wr \Sigma_{n+2}$. (See Example 12.) In particular, the realisation of $\mathcal{C}^n$ has the homotopy type of the wedge of $(2^n - 1)$ copies of the n -dimensional sphere. To obtain this description observe that the vertex set $H_{n+2}/H_{n+1} = (C_2 \wr \Sigma_{n+2})/(C_2 \wr \Sigma_{n+1})$ can be identified with $\{\pm 1, \dots, \pm(n+2)\}$ via the map sending the coset of $((\epsilon_1, \dots, \epsilon_{n+2}), \sigma)$ to $\epsilon_{n+2}\sigma(n+2)$, and that under this identification the k -simplex with lift $((\epsilon_1, \dots, \epsilon_{n+2}), \sigma)$ is

$$C = \{\epsilon_{n-k+2}\sigma(n-k+2), \dots, \epsilon_{n+2}\sigma(n+2)\},$$

so that a subset of $\{\pm 1, \dots, \pm(n+2)\}$ spans a simplex if and only if it does not contain any element and its negative.

Recall from Definition 3.4 of [11] that a simplicial complex is called *weakly Cohen-Macaulay of dimension n* if it is $(n-1)$ -connected and the link of each p -simplex is $(n-p-2)$ -connected. In each of the three examples above, $\mathcal{C}^n$ has the homotopy type of a wedge of n -dimensional spheres, and so is $(n-1)$ -connected. In fact, it is not hard to see that in these examples $\mathcal{C}^n$ is weakly Cohen-Macaulay of dimension n . This is an instance of the following general fact.

Theorem 14. $\mathcal{C}^n$ is weakly Cohen-Macaulay of dimension n .

The proof of this theorem is assembled from Propositions 15, 21 and 22, which are proved over the course of the next three sections.

Proof. By Proposition 15, if C is a p -simplex of $\mathcal{C}^n$ then $\text{lk}_{\mathcal{C}^n}(C) \cong \mathcal{C}^{n-p-1}$. It therefore suffices to show that $\mathcal{C}^n$ is $(n-1)$ -connected for all n , or equivalently that the barycentric subdivision $\text{sd } \mathcal{C}^n$ is $(n-1)$ -connected for all n . Now Proposition 21 shows that $|\text{sd } \mathcal{C}^n|$ is homeomorphic to the basic construction $\mathcal{U}(W_n, |\Delta|)$, while Proposition 22 shows that $\mathcal{U}(W_n, |\Delta|)$ is $(n-1)$ -connected. This completes the proof. $\square$

5. LINKS OF SIMPLICES OF $\mathcal{C}^n$

Proposition 15. Let C be a p -simplex of $\mathcal{C}^n$. Then $\text{lk}_{\mathcal{C}^n}(C) \cong \mathcal{C}^{n-p-1}$.

Proof. Choose a lift c of C . Define

$$\phi: W_{n-p-1}/W_{n-p-2} \longrightarrow W_n/W_{n-1}$$

by $\phi(dW_{n-p-2}) = cds_{n-p} \cdots s_n W_{n-1}$ for $d \in W_{n-p-1}$. This is well defined since every generator of W_{n-p-2} commutes with $s_{n-p}, \dots, s_n$. Observe that the domain and range of ϕ are the vertex sets of $\mathcal{C}^{n-p-1}$ and $\mathcal{C}^n$ respectively.

Claim 1. ϕ is an injection.

To prove this claim let $d, d' \in W_{n-p-1}$ satisfy

$$cd(s_{n-k} \cdots s_n)W_{n-1} = cd's_{n-k} \cdots s_n W_{n-1}.$$

Then

$$d^{-1}d' \in W_{n-p-1} \cap (s_{n-p} \cdots s_n)W_{n-1}(s_n \cdots s_{n-p}) = W_{n-p-2},$$

the latter equation by Proposition 6. Thus $d'W_{n-p-2} = dW_{n-p-2}$.

Claim 2. ϕ sends simplices of $\mathcal{C}^{n-p-1}$ to simplices of $\text{lk}_{\mathcal{C}^n}(C)$;

To prove this, suppose that D is an i -simplex of $\mathcal{C}^{n-p-1}$. Let $d \in W_{n-p-1}$ be a lift of D . Then

$$\phi D = \{cds_{n-p-i} \cdots s_n W_{n-1}, \dots, cds_{n-p} \cdots s_n W_{n-1}\}$$

while

$$\begin{aligned} C &= \{cs_{n-p+1} \cdots s_n W_{n-1}, \dots, cs_n W_{n-1}, cW_{n-1}\} \\ &= \{cds_{n-p+1} \cdots s_n W_{n-1}, \dots, cds_n W_{n-1}, cdW_{n-1}\}. \end{aligned}$$

Thus $\phi D \cap C = \emptyset$ by Proposition 8, and $\phi D \cup C$ is a simplex of $\mathcal{C}^n$ with lift cd , so that ϕD is a simplex of $\text{lk}_{\mathcal{C}^n}(C)$ as claimed.

Claim 3. Every simplex of $\text{lk}_{\mathcal{C}^n}(C)$ has the form ϕD for some simplex D of $\mathcal{C}^{n-p-1}$.

To prove this, suppose that $\bar{D}$ is an i -simplex of $\text{lk}_{\mathcal{C}^n}(C)$. Then $\bar{D} \cap C = \emptyset$ and $\bar{D} \cup C$ is a simplex of $\mathcal{C}^n$. Let c' be a lift of $\bar{D} \cup C$, and assume without loss that the ordering it induces on $\bar{D} \cup C$ contains $\bar{D}$ as an initial segment and C as a terminal segment with the ordering induced by c . Thus

$$\bar{D} = \{c'(s_{n-p-i} \cdots s_n)W_{n-1}, \dots, c'(s_{n-p} \cdots s_n)W_{n-1}\}$$

and

$$c'(s_{n-p+j} \cdots s_n)W_{n-1} = c(s_{n-p+j} \cdots s_n)W_{n-1}$$

for $j = 1, \dots, p+1$. The latter gives $c^{-1}c' \in W_{n-p-1}$ by Proposition 7, so that $c' = cd$ for some $d \in W_{n-p-1}$. Then $\bar{D} = \phi D$, where D is the i -simplex of $\mathcal{C}^{n-p-1}$ with lift d .

We can now prove the proposition. Combining the first claim with the third in the case of 0-simplices, we see that ϕ is an isomorphism between the vertex sets of $\mathcal{C}^{n-p-1}$ and $\text{lk}_{\mathcal{C}^n}(C)$. The second and third claims then show that ϕ induces an isomorphism of simplicial complexes from $\mathcal{C}^{n-p-1}$ to $\text{lk}_{\mathcal{C}^n}(C)$. $\square$

6. THE ACTION OF W_n ON $\mathcal{C}^n$ AND $\text{sd } \mathcal{C}^n$

Recall that if X is an abstract simplicial complex, then its *barycentric subdivision* $\text{sd } X$ is the abstract simplicial complex whose vertices are the simplices of X , and in which $\{C_0, \dots, C_k\}$ is a simplex if, after possibly reordering the elements, we have $C_0 \subset \dots \subset C_k$.

Here we will study the barycentric subdivision $\text{sd } \mathcal{C}^n$. The action of W_n on $\mathcal{C}^n$ induces an action of W_n on $\text{sd } \mathcal{C}^n$. This new action automatically has the property that if $w \in W_n$ and $\mathbf{C}$ is a simplex of $\text{sd } \mathcal{C}^n$, then w fixes every vertex of $\mathbf{C} \cap w\mathbf{C}$. (See p.115 of [3].) In particular any element of the stabilizer of $\mathbf{C}$ must fix $\mathbf{C}$ pointwise.

Lemma 16. *The action of W_n on $\mathcal{C}^n$ is transitive on the set of k -simplices for each k . For a k -simplex C of $\mathcal{C}^n$, any permutation of the vertices of C is realised by some element of W_n that fixes C . Every simplex of $\mathcal{C}^n$ is a face of an n -simplex.*

Proof. For the first claim, let C and D be k -simplices of $\mathcal{C}^n$ with respective lifts c and d . Then $(dc^{-1})C = D$. For the second claim, let C be a simplex of $\mathcal{C}^n$ with lift c , and let ϕ be a permutation of the vertices of C . Let c' be the lift of C whose induced ordering is obtained from the ordering induced by c by applying ϕ . Then $c'c^{-1}$ sends each vertex of C to its image under ϕ . The final claim follows because a k -simplex with lift c is a face of the n -simplex with lift c . $\square$

Lemma 17. *W_n acts transitively on the n -simplices of $\text{sd } \mathcal{C}^n$. Every simplex of $\text{sd } \mathcal{C}^n$ is a face of an n -simplex.*

Proof. Since $\mathcal{C}^n$ has dimension n , the set of n -simplices of $\text{sd } \mathcal{C}^n$ is in natural bijection with the set of n -simplices of $\mathcal{C}^n$ equipped with an ordering of their vertices. Lemma 16 shows that the action of W_n on this set is transitive, proving the first claim. The same lemma shows that every simplex of $\mathcal{C}^n$ is a face of an n -simplex, and the second claim follows immediately. $\square$


To understand the action of W_n on $\text{sd } \mathcal{C}^n$ we may now concentrate on a single n -simplex.

Definition 18. Let $\Delta = \{a_0, \dots, a_n\}$ denote the n -simplex of $\text{sd } \mathcal{C}^n$ with vertices $a_i = \{(s_{i+1} \dots s_n)W_{n-1}, \dots, s_n W_{n-1}, eW_{n-1}\}$ for $i = 1, \dots, n$. For each $s \in S_n$ define a face Δ_s of Δ as follows. If $s \in S_i \setminus S_{i-1}$ for $i = 1, \dots, n$ then


$$\Delta_s = \{a_0, \dots, \hat{a}_i, \dots, a_n\},$$

and if $s \in S_{-1}$ then $\Delta_s = \Delta$.

Example 19 (The case $n = 2$). Let $n = 2$. The diagram on the left shows the 2-simplex $\{s_1 s_2 W_1, s_2 W_1, W_1\}$ of $\mathcal{C}^2$ with lift the identity element, and the diagram on the right shows Δ as a face of the subdivision of this simplex.



The next diagrams show Δ_{s_2} , Δ_{s_1} , Δ_t and Δ_u for $t \in S_0 \setminus S_{-1}$, and $u \in S_{-1}$.



Observe that s_2 acts on $\{s_1s_2W_1, s_2W_1, W_1\}$ as the reflection that fixes $s_1s_2W_1$ and interchanges W_1 and s_2W_1 , and Δ_{s_2} is exactly the part of Δ fixed by this reflection. Similarly, s_1 acts as the reflection that fixes W_1 and interchanges $s_1s_2W_1$ and s_2W_1 , and Δ_{s_1} is precisely the part of Δ fixed by this reflection. Next, t can be thought of as “reflection in the edge $\{W_1, s_2W_1\}$ ”, and Δ_t is precisely the part of Δ that it fixes. Finally, u acts trivially on Δ since it lies in W_1 and commutes with s_1 and s_2 , and indeed $\Delta_u = \Delta$ is the part of Δ fixed by u .

Lemma 20. *Let F be a face of Δ . Then the stabilizer of F under the action of W_n is the subgroup generated by those $s \in S_n$ for which $F \subset \Delta_s$.*

Proof. For the purposes of this proof, given $i \geq 0$ we write $S_{=i}$ for the difference $S_i \setminus S_{i-1}$. So for $i \geq 1$ we have $S_{=i} = \{s_i\}$, while $S_{=0}$ is the set of elements of S_0 that do not commute with s_1 .

For $i \geq 0$, the stabilizer of a_i is the subgroup of W_n generated by $S_n \setminus S_{=i}$. For Proposition 5 shows that $s_{i+1}, \dots, s_n$ fix $a_i = \{s_{i+1} \cdots s_n W_{n-1}, \dots, s_n W_{n-1}, W_{n-1}\}$ and generate all permutations of its elements, while Proposition 7 shows that the subgroup consisting of elements that fix every element of a_i is W_{i-1} .

Let $F = \{a_{i_1}, \dots, a_{i_r}\}$. Then the stabilizer of F is the intersection of the stabilizers of the a_{i_j} . By the last paragraph this is the intersection of the subgroups generated by the sets $S_n \setminus S_{=i_j}$, and by a general result (see Theorem 4.1.6 of [5]) this is the subgroup generated by $\bigcap (S_n \setminus S_{=i_j}) = S_n \setminus \bigcup S_{=i_j}$.

Now $F \subset \Delta_s$ for all $s \in S_{-1}$, and $F \subset \Delta_s$ for $s \in S_{=i}$ if and only if $a_i \notin F$. Thus the set of s such that $F \subset \Delta_s$ is $S \setminus \bigcup S_{=i_j}$. Thus the subgroup generated by the s such that $F \subset \Delta_s$ is precisely the pointwise stabilizer of F . $\square$

7. THE BARYCENTRIC SUBDIVISION OF $\mathcal{C}^n$ AND THE BASIC CONSTRUCTION

The ‘basic construction’ is a method for building and studying certain spaces with group action. It can be used, for example, to study the topology of the Coxeter complex and Davis complex of a Coxeter group. In this section we will show that $|\text{sd } \mathcal{C}^n|$ is an instance of the basic construction, and we will use this to show that it is $(n-1)$ -connected.

To begin we recall the relevant notions from section 5.1 of [5]. (These are tailored to the case of Coxeter groups. For an approach to the basic construction that applies to more general groups see chapter II.12 of [4].) Let (W, S) be a Coxeter system. A *mirrored space* over S is a space X together with subspaces $X_s \subset X$, called *mirrors*, one for each $s \in S$. We assume that X is a CW-complex and that the mirrors are subcomplexes. The *basic construction* is the

space

$$\mathcal{U}(W, X) = (W \times X) / \sim$$

where $(v, x) \sim (w, y)$ if and only if $x = y$ and $v^{-1}w$ belongs to the subgroup generated by the $s \in S$ for which $x \in X_s$. The basic construction is equipped with the action of W by left translation, and we identify X with the image of $\{e\} \times X$ in $\mathcal{U}(W, X)$. Observe that $\mathcal{U}(W, X)$ has the structure of a CW-complex in which each translate wX is a subcomplex.

For us the most important feature of the basic construction is that it can be described as an increasing union of chambers, meaning copies of X , as we now recall from section 8.1 of [5]. Given $w \in W$, let $\text{In}(w) = \{s \in S \mid \ell(ws) < \ell(w)\}$ denote the set of letters with which a reduced expression for w can end, and let $X^{\text{In}(w)} = \bigcup_{s \in \text{In}(w)} X_s$ denote the corresponding union of mirrors. Now order the elements of W as $w_0, w_1, w_2, \dots$ where $w_0 = e$ and $\ell(w_m) \leq \ell(w_{m+1})$ for $m \geq 0$. Define

$$P_m = \bigcup_{i=0}^m w_i X,$$

so that $\mathcal{U}(W, X)$ is the increasing union of the subcomplexes P_m . Then

$$P_m = P_{m-1} \cup w_m X \quad \text{and} \quad P_{m-1} \cap w_m X = w_m X^{\text{In}(w_m)}.$$

The latter equation is by Lemma 8.1.1 of [5], and it will be useful to us since it specifies exactly how each chamber is attached to its predecessor.

Now let us show that $|\text{sd } \mathcal{C}^n|$ is an instance of the basic construction. Fix the Coxeter system (W_n, S_n) . Recall the simplex Δ and the faces Δ_s introduced in Definition 18. We make $|\Delta|$ into a mirrored space over S_n by defining the mirror $|\Delta|_s$ to be the subspace $|\Delta_s|$ of $|\Delta|$. We may therefore form the basic construction $\mathcal{U}(W_n, |\Delta|)$. The inclusion $|\Delta| \hookrightarrow |\text{sd } \mathcal{C}^n|$ sends $|\Delta|_s$ into the subset fixed by s , and so extends uniquely to a W_n -equivariant map $\mathcal{U}(W_n, |\Delta|) \rightarrow |\text{sd } \mathcal{C}^n|$.

Proposition 21. *The map $\mathcal{U}(W_n, |\Delta|) \rightarrow |\text{sd } \mathcal{C}^n|$ is a homeomorphism.*

Proof. The map is surjective because any point of $|\text{sd } \mathcal{C}^n|$ is in a translate of $|\Delta|$. This follows from Lemma 17, which shows that every simplex of $\text{sd } \mathcal{C}^n$ is a face of a translate of Δ .

The map is injective because if $x \in |\Delta|$, then the image of x in $\mathcal{U}(W_n, |\Delta|)$ has stabilizer generated by those s for which $x \in |\Delta|_s$. To see this, let F denote the unique face of Δ for which x lies in the interior of $|F|$. Then $x \in |\Delta|_s = |\Delta_s|$ if and only if $F \subset \Delta_s$, and the stabilizer of x is precisely the stabilizer of F . The result then follows from Lemma 20.

The map is a homeomorphism because $|\text{sd } \mathcal{C}^n|$ has the weak topology with respect to the realizations of its simplices. By Lemma 17 this coincides with the weak topology with respect to the realizations of its n -simplices. This is exactly the topology of $\mathcal{U}(W_n, |\Delta|)$. $\square$

Proposition 22. *$\mathcal{U}(W_n, |\Delta|)$ is $(n-1)$ -connected.*

This relies on the following two lemmas.

Lemma 23. *For $w \in W_n$, $w \neq e$, the space $|\Delta|^{\text{In}(w)}$ is $(n-2)$ -connected.*

Proof. The set $\text{In}(w)$ is nonempty since $w \neq e$. Thus $|\Delta|^{\text{In}(w)}$ is either $|\Delta|$, or it is a nonempty union of facets of $|\Delta|$. In the first case it is contractible, and in the second case it is either contractible (if not all facets are in the union) or it is $\partial|\Delta| \cong S^{n-1}$ (if all facets are in the union). In all cases it is $(n-2)$ -connected. $\square$

Lemma 24. *Let $n \geq 1$. Suppose that $(X; A, B)$ is a CW-triad in which A and B are $(n-1)$ -connected and $C = A \cap B$ is $(n-2)$ -connected. Then X is $(n-1)$ -connected.*

Proof. For $n = 1$ this is immediate since the union of two path-connected spaces with nonempty intersection is path-connected. So we assume that $n \geq 2$. The pairs (A, C) and (B, C) are $(n-1)$ -connected, and C is path-connected, so that Theorem 4.23 of [9] can be applied to show that $\pi_i(A, C) \rightarrow \pi_i(X, B)$ is an isomorphism for $i < 2n-2$, and in particular for $i \leq (n-1)$. Thus (X, B) is $(n-1)$ -connected, and the same then follows for X itself. $\square$

Proof of Proposition 22. If $n = 0$ then the claim is that $\mathcal{U}(W_n, |\Delta|)$ is nonempty, which holds vacuously. So we may assume that $n \geq 1$.

As in the discussion at the start of the section, order the elements of W_n as $w_0, w_1, w_2, \dots$ starting with the identity and respecting the length. Then $\mathcal{U}(W_n, |\Delta|)$ is the union of subcomplexes $P_0 \subset P_1 \subset P_2 \subset \dots$ where $P_0 = |\Delta|$ and

$$P_m = P_{m-1} \cup w_m |\Delta| \quad \text{with} \quad P_{m-1} \cap w_m |\Delta| = w_m |\Delta|^{\text{In}(w_m)}.$$

It will suffice to show that each P_m is $(n-1)$ -connected. We do this by induction on m .

In the initial case $m = 0$ we have $P_0 = e|\Delta|$, which is contractible and so the claim holds. For the induction step we take $m \geq 1$ and assume that P_{m-1} is $(n-1)$ -connected. Then $P_m = P_{m-1} \cup w_m |\Delta|$ is the union of the subcomplexes P_{m-1} and $w_m |\Delta|$, and their intersection $w_m |\Delta|^{\text{In}(w_m)}$ is $(n-2)$ -connected by Lemma 23. Thus $(P_m; P_{m-1}, w_m |\Delta|)$ is a CW-triad in which the subspaces P_{m-1} and $w_m |\Delta|$ are $(n-1)$ -connected and their intersection is $(n-2)$ -connected. It now follows from Lemma 24 that P_m is $(n-1)$ -connected as required. $\square$

8. THE ORDERED SIMPLICES OF $\mathcal{C}^n$

In this section we introduce a semisimplicial set $\mathcal{D}^n$ and identify it as the semisimplicial set of ordered simplices in $\mathcal{C}^n$. We then use the fact that $\mathcal{C}^n$ is weakly Cohen-Macaulay of dimension n to deduce that the geometric realisation $\|\mathcal{D}^n\|$ is $(n-1)$ -connected, an approach we learned from Wahl's paper [17] (see in particular Proposition 7.9 of [17], which is due to Randal-Williams). In this section and the next we will use semisimplicial spaces and their realisations. The background material we require can be found in section 2 of [15].

Definition 25. Let $\mathcal{D}^n$ denote the semisimplicial set with k -simplices.

$$\mathcal{D}_k^n = \begin{cases} W_n / W_{n-k-1} & k \leq n \\ \emptyset & k > n \end{cases}$$

and with face maps

$$d_i: W_n / W_{n-k-1} \longrightarrow W_n / W_{n-k}$$

defined by

$$d_i(cW_{n-k-1}) = c(s_{n-k+i} \cdots s_{n-k+1})W_{n-k}$$

for $i = 0, \dots, k$.

It is a simple exercise to verify that the face maps d_i satisfy the relations $d_i \circ d_j = d_{j+1} \circ d_i$ for $i < j$. Alternatively, it is a consequence of the proof of Proposition 27 below.

Definition 26. Let X be a simplicial complex. By an *ordered simplex* of X , we mean a simplex of X equipped with an ordering of its vertices. The *semi-simplicial set of ordered simplices in X* , denoted X^{ord} , has for its k -simplices the ordered k -simplices in X , with face maps d_i given by forgetting the i -th vertex of an ordered simplex.

Proposition 27. $\mathcal{D}^n$ is isomorphic to $\mathcal{C}^{n, \text{ord}}$.

Proof. We define $\phi_k: \mathcal{D}_k^n \rightarrow \mathcal{C}_k^{n, \text{ord}}$ by

$$\phi_k(cW_{n-k-1}) = \{c(s_{n-k+1} \cdots s_n)W_{n-1}, \dots, cs_n W_{n-1}, cW_{n-1}\}$$

for $cW_{n-k-1} \in W_n/W_{n-k-1}$. In other words, $\phi_k(cW_{n-k-1})$ is the k -simplex with lift c , equipped with the ordering induced by c . The map ϕ_k is well defined because the generators of W_{n-k-1} all commute with $s_{n-k+1}, \dots, s_n$. It is surjective because by definition every simplex admits a lift, and any ordering of a simplex is afforded by some lift (see the paragraph following Definition 9). It is injective because if $\phi_k(cW_{n-k-1}) = \phi_k(c'W_{n-k-1})$ then $cs_i \cdots s_n W_{n-1} = c's_i \cdots s_n W_{n-1}$ for $i = n - k + 1, \dots, n + 1$, so that $cW_{n-k-1} = c'W_{n-k-1}$ by Proposition 7.

To complete the proof we must show that the face maps in $\mathcal{C}^{n, \text{ord}}$ and $\mathcal{D}^n$ are compatible under the ϕ_k . In other words, given $0 \leq i \leq k \leq n$, we must show that

$$\phi_{k-1} \circ d_i = d_i \circ \phi_k.$$

Observe from the definition of d_i in $\mathcal{D}^n$ that for $i \geq 1$ we have $d_i(cW_{n-k-1}) = d_{i-1}(cs_{n-k+i}W_{n-k+1})$. On the other hand, Proposition 5 shows that $\phi_k(cW_{n-k-1})$ and $\phi_k(cs_{n-k+i}W_{n-k+1})$ differ only by the transposition of their $(i-1)$ -st and i -th vertices, so that $d_i(\phi_k(cW_{n-k-1})) = d_{i-1}(\phi_k(cs_{n-k+i}W_{n-k+1}))$. Thus the claim will follow by induction on i so long as we can show that

$$\phi_{k-1} \circ d_0 = d_0 \circ \phi_k.$$

This follows by inspection. □

Corollary 28. $\|\mathcal{D}^n\|$ is $(n-1)$ -connected.

Proof. Theorem 14 shows that $\mathcal{C}^n$ is weakly Cohen-Macaulay of dimension n . Proposition 7.9 of [17] shows that if a simplicial complex X is weakly Cohen-Macaulay of dimension n , then $\|X^{\text{ord}}\|$ is $(n-1)$ -connected. Consequently $\|\mathcal{C}^{n, \text{ord}}\|$ is $(n-1)$ -connected, and by Proposition 27 the same holds for $\|\mathcal{D}^n\|$. □

9. COMPLETION OF THE PROOF

We now complete the proof of the main theorem. This section is modelled closely on section 5 of [15], from which there is little essential difference. It is also similar to the proof of Theorem 2 of [12].

We regard $\mathcal{D}^n$ as a simplicial space by equipping its constituent sets with the discrete topology. Then we form a semisimplicial space

$$EW_n \times_{W_n} \mathcal{D}^n$$

by setting $(EW_n \times_{W_n} \mathcal{D}^n)_k = EW_n \times_{W_n} \mathcal{D}_k^n$ and using the face maps obtained from those of $\mathcal{D}^n$.

Lemma 29. *The projection $EW_n \times_{W_n} \mathcal{D}_0^n \rightarrow BW_n$ makes $EW_n \times_{W_n} \mathcal{D}^n$ into an augmented simplicial space over BW_n , and the induced map $\|EW_n \times_{W_n} \mathcal{D}^n\| \rightarrow BW_n$ is $(n-1)$ -connected.*

Proof. The composites of the projection with d_0 and d_1 coincide, so that the projection is indeed an augmentation. Since $EW_n \rightarrow BW_n$ is a locally trivial principal W_n -bundle, it follows that $\|EW_n \times_{W_n} \mathcal{D}^n\| \rightarrow BW_n$ is a locally trivial bundle with fibre $\|W_n \times_{W_n} \mathcal{D}^n\| \cong \|\mathcal{D}^n\|$, which is $(n-1)$ -connected by Corollary 28, so that the map itself is $(n-1)$ -connected. $\square$

Lemma 30. *There are homotopy equivalences $EW_n \times_{W_n} \mathcal{D}_k^n \simeq BW_{n-k-1}$ under which the face maps $d_i: EW_n \times_{W_n} \mathcal{D}_k^n \rightarrow EW_n \times_{W_n} \mathcal{D}_{k-1}^n$ are all homotopic to the stabilization map $BW_{n-k-1} \rightarrow BW_{n-k}$, and under which the composite $EW_n \times_{W_n} \mathcal{D}_0^n \rightarrow \|EW_n \times_{W_n} \mathcal{D}^n\| \rightarrow BW_n$ is homotopic to the stabilization map $BW_{n-1} \rightarrow BW_n$.*

Proof. There is an isomorphism

$$EW_n \times_{W_n} \mathcal{D}_k^n = EW_n \times_{W_n} (W_n/W_{n-k-1}) \xrightarrow{\cong} EW_n/W_{n-k-1}$$

sending the orbit of (x, cW_{n-k-1}) to the orbit of $c^{-1}x$. This identifies d_i with the map

$$EW_n/W_{n-k-1} \longrightarrow EW_n/W_{n-k}$$

sending the W_{n-k-1} -orbit of x to the W_{n-k} -orbit of $(s_{n-k+1} \cdots s_{n-k+i})x$. Since $(s_{n-k+1} \cdots s_{n-k+i})$ commutes with every element of W_{n-k-1} , this map is homotopic to the one sending the W_{n-k-1} -orbit of x to the W_{n-k} -orbit of x . Now the equivariant homotopy equivalences

$$EW_{n-k-1} \rightarrow EW_n, \quad EW_{n-k} \rightarrow EW_n$$

induce homotopy equivalences

$$BW_{n-k-1} \rightarrow EW_n/W_{n-k-1}, \quad BW_{n-k} \rightarrow EW_n/W_{n-k}$$

under which the map $EW_n/W_{n-k-1} \rightarrow EW_n/W_{n-k}$ just described becomes the stabilization map. $\square$

The skeletal filtration of $\|EW_n \times_{W_n} \mathcal{D}^n\|$ leads to a first-quadrant spectral sequence

$$E_{k,l}^1 = H_l(EW_n \times_{W_n} \mathcal{D}_k^n) \implies H_{k+l}(\|EW_n \times_{W_n} \mathcal{D}^n\|)$$

in which the differential d^1 is given by the alternating sum $\sum_{i=0}^k (-1)^i (d_i)_*$ of the maps induced by the face maps. Lemma 30 allows us to identify the E^1 -term of this spectral sequence: $E_{k,l}^1 = H_l(BW_{n-k-1})$, and $d^1: E_{k,l}^1 \rightarrow E_{k-1,l}^1$ is the stabilization map $H_l(BW_{n-k-1}) \rightarrow H_l(BW_{n-k})$ if k is even, and is zero if k is odd.

Lemma 31. *Assume that for all $m < n$ the stabilization map $H_l(BW_{m-1}) \rightarrow H_l(BW_m)$ is an isomorphism in degrees $2l \leq m$. Then the spectral sequence has the following properties:*

- (1) $E_{0,l}^\infty = \cdots = E_{0,l}^2 = E_{0,l}^1$ for $2l \leq n$.
- (2) $E_{k,l}^\infty = 0$ for $k > 0$ and $2(k+l) \leq n$.

Proof. The assumption allows us to deduce that $E_{k,l}^2 = 0$ when $k \geq 1$ is odd and $2l + k + 1 \leq n$, and that $E_{k,l}^2 = 0$ when $k \geq 2$ is even and $2l + k \leq n$. For in the first case $d^1: E_{k+1,l}^1 \rightarrow E_{k,l}^1$ is the stabilization map $H_l(BW_{n-k-2}) \rightarrow H_l(BW_{n-k-1})$, and in the second case $d^1: E_{k,l}^1 \rightarrow E_{k-1,l}^1$ is the stabilization map $H_l(BW_{n-k-1}) \rightarrow H_l(BW_{n-k})$, and our assumption means that both are isomorphisms in the given range.

To prove the first claim, observe that $E_{0,l}^2 = E_{0,l}^1$ since $d^1: E_{1,l}^1 \rightarrow E_{0,l}^1$ is zero. The remaining differentials with target in bidegree $(0, l)$ are $d^k: E_{k,l-k+1}^k \rightarrow E_{0,l}^k$ with $k \geq 2$, and these have domain zero since $2(l-k+1)+k \leq 2(l-k+1)+k+1 = 2l - k + 2 \leq 2l \leq n$ so that $E_{k,l-k+1}^k = 0$. To prove the second claim, observe that if $2(k+l) \leq n$ and $k > 0$, then certainly $2l + k < 2l + k + 1 \leq 2(l+k) \leq n$, so that $E_{k,l}^2 = 0$. $\square$

We can now complete the proof of the main theorem, showing by induction on $n \geq 2$ that $H_l(BW_{n-1}) \rightarrow H_l(BW_n)$ is an isomorphism for $2l \leq n$. For $n = 2$ the claim is that $H_l(BW_1) \rightarrow H_l(BW_2)$ is an isomorphism for $l = 0, 1$. For $l = 0$ this is trivial since both BW_1 and BW_2 are connected. For $l = 1$ this follows from the well-known fact that if (W, S) is a Coxeter system then $H_1(BW)$ is isomorphic to the elementary abelian 2-group generated by the elements of S , subject to the relation that identifies $s, t \in S$ if m_{st} is odd. Take $n > 2$ and suppose that the theorem holds for all smaller integers. Lemma 30 shows that the composite

$$H_l(W_{n-1}) = E_{0,l}^1 \rightarrow E_{0,l}^\infty \rightarrow H_l(\|EW_n \times_{W_n} \mathcal{D}^n\|) \rightarrow H_l(BW_n)$$

is the stabilization map, and we must show that this is an isomorphism for $2l \leq n$. Lemma 31 shows that the first two arrows are isomorphisms in this range, while Lemma 29 shows that the last map is an isomorphism for $l \leq n-1$, which holds since $2l \leq n$ and $n \geq 3$.

REFERENCES

- [1] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [2] Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [3] Glen E. Bredon. *Introduction to compact transformation groups*. Academic Press, New York-London, 1972. Pure and Applied Mathematics, Vol. 46.

- [4] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [5] Michael W. Davis. *The geometry and topology of Coxeter groups*, volume 32 of *London Mathematical Society Monographs Series*. Princeton University Press, Princeton, NJ, 2008.
- [6] Søren Galatius and Oscar Randal-Williams. Homological stability for moduli spaces of high dimensional manifolds. I. [arXiv:1403.2334v1](#), 2014.
- [7] John L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [8] Allen Hatcher. Homological stability for automorphism groups of free groups. *Comment. Math. Helv.*, 70(1):39–62, 1995.
- [9] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [10] Allen Hatcher and Karen Vogtmann. Homology stability for outer automorphism groups of free groups. *Algebr. Geom. Topol.*, 4:1253–1272, 2004.
- [11] Allen Hatcher and Nathalie Wahl. Stabilization for mapping class groups of 3-manifolds. *Duke Math. J.*, 155(2):205–269, 2010.
- [12] Moritz C. Kerz. The complex of words and Nakaoka stability. *Homology Homotopy Appl.*, 7(1):77–85, 2005.
- [13] Minoru Nakaoka. Decomposition theorem for homology groups of symmetric groups. *Ann. of Math. (2)*, 71:16–42, 1960.
- [14] Daniel Quillen. Finite generation of the groups K_i of rings of algebraic integers. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 179–198. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [15] Oscar Randal-Williams. Homological stability for unordered configuration spaces. *Q. J. Math.*, 64(1):303–326, 2013.
- [16] Louis Solomon. A Mackey formula in the group ring of a Coxeter group. *J. Algebra*, 41(2):255–264, 1976.
- [17] Nathalie Wahl. Homological stability for automorphism groups. [arXiv:1409.3541v1](#), 2014.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE,
UNITED KINGDOM

E-mail address: r.hepworth@abdn.ac.uk